

Datasheet for ABIN6938912

Mesothelin Protein (MSLN) (AA 296-580) (His tag)

2 Images



Overview

Overview	
Quantity:	100 μg
Target:	Mesothelin (MSLN)
Protein Characteristics:	AA 296-580
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Mesothelin protein is labelled with His tag.
Product Details	
Sequence:	AA 296-580
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 0.01 EU per µg by the LAL method.
Target Details	
Target:	Mesothelin (MSLN)
Alternative Name:	Mesothelin (MSLN Products)
Background:	Mesothelin (MSLN) is also known as CAK1 antigen, Pre-pro-megakaryocyte-potentiating factor,
	which belongs to the mesothelin family. Mesothelin / MSLN can be proteolytically cleaved into the following two chains by a furin-like convertase: Megakaryocyte-potentiating factor (MPF)

and the cleaved form of mesothelin. Both MPF and the cleaved form of mesothelin are N-

Target Details

glycosylated. Mesothelin / MSLN can interacts with MUC16. The membrane-anchored forms of	
MSLN may play a role in cellular adhesion. MPF potentiates megakaryocyte colony formation in	
vitro.	

Molecular Weight:

33.0 kDa

Pathways:

EGFR Signaling Pathway, Positive Regulation of Peptide Hormone Secretion, Intracellular
Steroid Hormone Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway,
Carbohydrate Homeostasis, cAMP Metabolic Process, Regulation of G-Protein Coupled
Receptor Protein Signaling, Positive Regulation of Endopeptidase Activity, Regulation of
Carbohydrate Metabolic Process

Application Details


Restrictions:

For Research Use only

Handling


Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C

Images



SDS-PAGE

Image 1. Human Mesothelin (296-580), His Tag, low endotoxin on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than $95\,\%$.



ELISA

Image 2. Immobilized Human Mesothelin (296-580), His Tag, low endotoxin (ABIN5954925,ABIN6938912) at $0.5~\mu$ g/mL (100 μ L/well) can bind A MAb, Human IgG1 with a linear range of 0.2-3 ng/mL (QC tested).