

Datasheet for ABIN6939024

**Recombinant anti-Cadherin-16 antibody****6** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	Cadherin-16 (CDH16)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Cadherin-16 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

## Product Details

Immunogen:	Recombinant human full-length CDH16 protein
Clone:	CDH16-1532R
Isotype:	IgG
Specificity:	This MAb recognizes a protein of 130 kDa, identified as Ksp-cadherin. Cadherins form a superfamily of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins have been implicated in embryogenesis, tissue morphogenesis, tissue structure maintenance, cell polarization, neoplastic invasiveness and metastasis, and membrane transport. It is suggested that Ksp-cadherin is a marker for terminal differentiation of the basolateral membranes of renal tubular epithelial cells. Within the kidney, Ksp-Cadherin is found exclusively in the basolateral

## Product Details

membrane of renal tubular epithelial cells and collecting duct cells, and not in glomeruli, renal interstitial cells, or blood vessels. Ksp-Cadherin has been suggested to distinguish Chromophobe Renal-Cell Carcinoma from Oncocytoma.

Cross-Reactivity (Details): Human, Mouse, Rat,

Purification: 200ug/ml of Ab Purified by Protein A.

## Target Details

Target: Cadherin-16 (CDH16)

Alternative Name: CDH16 ([CDH16 Products](#))

Background: Cadherin-16 (CDH16), Kidney-specific cadherin, Ksp-cadherin antibody, Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16  
Cellular localisation: Cell Surface and Cytoplasmic

Molecular Weight: 130kDa

Gene ID: 1014, 513660

UniProt: [O75309](#)

## Application Details

Application Notes: Positive Control: Normal kidney or renal cell carcinoma.  
Known Application: Western Blot (1-2 µg/mL), Flow Cytometry (1-2 µg/million cells), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA buffer, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Also available WITHOUT BSA & azide at 1.0mg/ml.
Expiry Date:	24 months

Images




Western Blotting

**Image 1.** Western Blot Analysis of human Kidney lysate using KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)

Flow Cytometry

**Image 2.** Flow Cytometric Analysis of Human HEK293 cells using KSP-Cadherin Recombinant Rabbit Monoclonal Antibody (CDH16/1532R) followed by Goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).



#### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN6939024.