
# antibodies -online.com







# anti-CDX2 antibody (AA 150-249)





_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μg	
Target:	CDX2	
Binding Specificity:	AA 150-249	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CDX2 antibody is un-conjugated	
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Immunostaining (ISt)	

## **Product Details**

Immunogen:	Recombinant fragment (around aa150-249) of human CDX2 protein (exact sequence is proprietary)
Clone:	CDX2-1690
Isotype:	lgG2a
Specificity:	The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CDX2 has been useful to establish GI origin of metastatic adenocarcinomas and carcinoidsand is especially useful to distinguish metastatic colorectal adenocarcinoma from lung adenocarcinoma. However, mucinous carcinomas of the ovary also express CDX2 protein. It limits the usefulness of this marker in the


## **Product Details**

- Todaot Dotano		
	distinction of metastatic colorectal adenocarcinoma from mucinous carcinoma of the ovary.	
Cross-Reactivity (Details):	Human,	
Purification:	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G.	
Target Details		
Target:	CDX2	
Alternative Name:	CDX2 (CDX2 Products)	
Background:	Caudal type homeobox 2, Caudal type homeobox transcription factor 2, Caudal-type homeobo protein 2, CDX2,CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker)  Cellular localisation: Nuclear	
Molecular Weight:	40kDa	
Gene ID:	1045, 174249	
UniProt:	Q99626	
Pathways:	Peptide Hormone Metabolism, Stem Cell Maintenance	
Application Details		
Application Notes:	Positive Control: HT29 cells. Colon Carcinoma.	
	Known Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA),	
	,Immunohistochemistry (Formalin-fixed) (1-2 $\mu$ g/mL for 30 min at RT),(Staining of formalin-	
	fixed tissues is enhanced by heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 fo	
	45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilution for a specific	
	application should be determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	

#### Handling


Storage:	4 °C,-80 °C	
Storage Comment:	Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Als available WITHOUT BSA & azide at 1.0mg/ml.	
Expiry Date:	24 months	

#### **Images**




#### **SDS-PAGE**

**Image 1.** SDS-PAGE Analysis Purified CDX2 Mouse Monoclonal Antibody (CDX2/1690). Confirmation of Purity and Integrity of Antibody.



#### **Protein Array**

Image 2. Analysis of Protein Array containing >19,000 fulllength human proteins using CDX2 Mouse Monoclonal Antibody (CDX2/1690) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Zscore, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb



to protein X is equal to 29.

## **Immunohistochemistry**

**Image 3.** Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CDX2 Mouse Monoclonal Antibody (CDX2/1690).

Please check the product details page for more images. Overall 4 images are available for ABIN6939057.