

Datasheet for ABIN6939094

Recombinant anti-CFTR antibody**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CFTR
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Application:	Immunostaining (ISt), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant full-length human CFTR protein.
Clone:	RCFTR-1342
Isotype:	IgG1
Specificity:	Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

Product Details

Cross-Reactivity (Details):	Human, Mouse,
Purification:	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G.

Target Details

Target:	CFTR
Alternative Name:	CFTR (CFTR Products)
Background:	ABC35, ATP Binding Cassette Superfamily C Member 7 (ABCC7), cAMP-dependent chloride channel, CFTR, CFTR/MRP, Channel conductance-controlling ATPase, Cystic Fibrosis Transmembrane Conductance Regulator, MRP7, TNR CFTR,CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Cellular localisation: Cell Surface and Cytoplasmic
Molecular Weight:	165-170kDa
Gene ID:	1080, 489786, 621460
UniProt:	P13569

Application Details

Application Notes:	Positive Control: MOLT-4 cells. Pancreas, Kidney or Placenta. Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

Handling


Concentration:	200 µg/mL
Buffer:	Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C

Handling

Storage Comment: Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Also available WITHOUT BSA & azide at 1.0mg/ml.


Expiry Date: 24 months

Images



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified CFTR Mouse Recombinant Monoclonal Antibody (rCFTR/1342). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Pancreas stained with CFTR Mouse Recombinant Monoclonal Antibody (rCFTR/1342).