# antibodies - online.com







## anti-UPK1A antibody (AA 114-173)

100 μg





_							
0	V	е	r١	/1	е	V	1

Quantity:

Target:	UPK1A
Binding Specificity:	AA 114-173
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))
Product Details	
Immunogen:	Recombinant fragment (around aa 114-173) of human Uroplakin 1A (UPK1A) protein (exact
	sequence is proprietary)
Clone:	UPK1A-2925
Isotype:	IgG1
Isotype: Specificity:	IgG1  The protein encoded by this gene is a member of the transmembrane 4 superfamily, also
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is found in the asymmetrical unit membrane (AUM) where it can
	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is found in the asymmetrical unit membrane (AUM) where it can complex with other transmembrane 4 superfamily proteins. It may play a role in normal bladder


#### **Product Details**

Product Details		
Cross-Reactivity (Details):	Human,	
Purification:	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G.	
Towns Dataile		
Target Details		
Target:	UPK1A	
Alternative Name:	UPK1A (UPK1A Products)	
Background:	UP1A, UPIA, UPKA, TSPAN21, Uroplakin 1A (Urothelial Differentiation Marker)	
	Cellular localisation: Cell Surface	
Molecular Weight:	29kDa	
Gene ID:	11045, 159309	
UniProt:	000322	
Application Details		
Application Notes:	Positive Control: Urinary Bladder.	
	Known Application: ELISA (For coating, order Ab without BSA), Immunohistochemistry	
	(Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires	
	heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95&degC followed	
	by cooling at RT for 20 minutes), Optimal dilution for a specific application should be	
	determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-80 °C	
Storage Comment:	Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Also	
	available WITHOUT BSA & azide at 1.0mg/ml.	

**Expiry Date:** 


24 months

#### **Images**




#### **SDS-PAGE**

**Image 1.** SDS-PAGE Analysis Purified Uroplakin 1A Mouse Monoclonal Antibody (UPK1A/2925). Confirmation of Purity and Integrity of Antibody.



#### **Protein Array**

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using Uroplakin 1A Mouse Monoclonal Antibody (UPK1A/2925) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



### **Immunohistochemistry**

Image 3. Formalin-fixed, paraffin-embedded human Urothelial Carcinoma stained with Uroplakin 1A Mouse Monoclonal Antibody (UPK1A/2925).

Please check the product details page for more images. Overall 4 images are available for ABIN6939118.