

Datasheet for ABIN6939124

**Recombinant anti-Chromogranin A antibody**[Go to Product page](#)**3** Images

## Overview

Quantity:	100 µg
Target:	Chromogranin A (CHGA)
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Chromogranin A antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant full-length human CHGA protein.
Clone:	RCHGA-413
Isotype:	IgG2b kappa
Specificity:	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, paragangliomas, and other neuroendocrine tumors. Co-expression of chromogranin A and neuron specific enolase (NSE) is common in neuroendocrine neoplasms. Reportedly, co-expression of certain keratins and chromogranin indicates neuroendocrine lineage. The presence of strong anti-chromogranin staining and absence of anti-keratin staining should raise the possibility of paraganglioma. The co-expression of chromogranin and NSE is typical of neuroendocrine neoplasms. Most pituitary

## Product Details

	adenomas and prolactinomas readily express chromogranin.
No Cross-Reactivity:	Rat (Rattus)
Purification:	Purified by Protein A/G

## Target Details

Target:	Chromogranin A (CHGA)
Alternative Name:	CHGA ( <a href="#">CHGA Products</a> )
Molecular Weight:	68-75kDa
Gene ID:	1113
UniProt:	<a href="#">P10645</a>
Pathways:	<a href="#">Negative Regulation of Hormone Secretion</a> , <a href="#">cAMP Metabolic Process</a> , <a href="#">Regulation of G-Protein Coupled Receptor Protein Signaling</a>


## Application Details

Application Notes:	Positive Control: PC12 cells. Adrenal gland, bowel, thyroid, pancreas or pheochromocytoma. Known Application: Immunohistochemistry (Formalin-fixed) (0.25-0.5 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes) Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

## Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months




SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified Chromogranin A Mouse Recombinant Monoclonal Ab (rCHGA/413). Confirmation of Purity and Integrity of Antibody.

Protein Array

**Image 2.** Analysis of Protein Array containing >19,000 full-length human proteins using Chromogranin A Recombinant Mouse Monoclonal Antibody (rCHGA/413) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Pancreas stained with Chromogranin A Mouse Recombinant Monoclonal Ab (rCHGA/413).