.-online.com antibodies

Datasheet for ABIN6939136 Recombinant anti-Chromogranin A antibody

4 Images



Overview

Quantity:	100 µg
Target:	Chromogranin A (CHGA)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Chromogranin A antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant human full-length chromogranin A(CHGA) protein
Clone:	CHGA-1731R
lsotype:	lgG
Specificity:	Chromogranin A is present in neuroendocrine cells throughout the body, including the
	neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is
	an excellent marker for carcinoid tumors, pheochromocytomas, paragangliomas, and other
	neuroendocrine tumors. Co-expression of chromogranin A and neuron specific enolase (NSE) is
	common in neuroendocrine neoplasms. Reportedly, co-expression of certain keratins and
	chromogranin indicates neuroendocrine lineage. The presence of strong anti-chromogranin
	staining and absence of anti-keratin staining should raise the possibility of paraganglioma. The
	co-expression of chromogranin and NSE is typical of neuroendocrine neoplasms. Most pituitary

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6939136 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Product Details	
	adenomas and prolactinomas readily express chromogranin.
No Cross-Reactivity:	Rat (Rattus)
Purification:	Purified by Protein A/G
Target Details	
Target:	Chromogranin A (CHGA)
Alternative Name:	CHGA (CHGA Products)
Molecular Weight:	68-75kDa
Gene ID:	1113
UniProt:	P10645
Pathways:	Negative Regulation of Hormone Secretion, cAMP Metabolic Process, Regulation of G-Protein Coupled Receptor Protein Signaling
Application Details	
Application Notes:	Positive Control: PC12 cells. Adrenal gland, bowel, parathyroid, pancreas or pheochromocytoma. Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1.0 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes) Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6939136 | 09/10/2023 | Copyright antibodies-online. All rights reserved.


Handling

is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date:


24 months

Images



SDS-PAGE


Image 1. SDS-PAGE Analysis of Purified Chromogranin A Rabbit Recombinant Monoclonal Antib (CHGA/1731R). Confirmation of Purity and Integrity.



Protein Array

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using Chromogranin A Rabbit Recombinant Monoclonal Antibody (CHGA/1731R). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. Sscore therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Images



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Pancreas stained with Chromogranin A Rabbit Recombinant Monoclonal Ab (CHGA/1731R).

Please check the product details page for more images. Overall 4 images are available for ABIN6939136.