antibodies .- online.com





anti-CNN1 antibody

2 Images



Overview

Quantity:	100 μg
Target:	CNN1
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CNN1 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC), Staining Methods (StM)

Product Details


Immunogen:	Recombinant human CNN1 protein
Isotype:	IgG
Specificity:	Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist, a 34 kDa form and a 29 kDa form. Expression of the 29 kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this antibody reacts with only the 34 kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.

Product Details Purified by Protein A/G Purification: Target Details Target: CNN1 CNN1 (CNN1 Products) Alternative Name: Molecular Weight: 34kDa Gene ID: 1264 UniProt: P51911 **Application Details** Positive Control: Myoepithelial cells in breast ducts. **Application Notes:** Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined. Restrictions: For Research Use only Handling Concentration: 200 μg/mL 10 mM PBS with 0.05 % BSA & 0.05 % azide. Buffer: Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. Storage: 4 °C.-80 °C Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


24 months

Expiry Date:



Immunohistochemistry

Image 1. Formalin-fixed, paraffin-embedded Rat Uterus stained with Calponin Rabbit Polyclonal Antibody



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Uterus stained with Calponin Rabbit Polyclonal Antibody