

Datasheet for ABIN6939194

**Recombinant anti-G-CSF antibody****2** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	G-CSF (CSF3)
Reactivity:	Human, Macaque
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This G-CSF antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant full-length human CSF3 protein
Clone:	CSF3-3166R
Isotype:	IgG
Specificity:	This MAb recognizes granulocyte-colony stimulating factor (G-CSF) in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell types. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of myeloid cells. It is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes. G-CSF is a pleiotropic cytokine that influences differentiation, proliferation and activation of the neutrophilic granulocyte lineage. The human G-CSF cDNA encodes a 207 amino acid precursor containing a 29 amino acid signal peptide that is

## Product Details

proteolytically cleaved to form a 178 amino acid residue mature protein. Two G-CSF s, which are identical except for a three amino acid deletion in the amino-terminus of one form of the protein have been isolated from human cells. Murine and human G-CSF s share 73 % sequence identity at the amino acid level.

Purification: Purified by Protein A/G

## Target Details

Target: G-CSF (CSF3)

Alternative Name: CSF3 ([CSF3 Products](#))

Molecular Weight: 19kDa

Gene ID: 1440

UniProt: [P09919](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Regulation of Actin Filament Polymerization](#)

## Application Details

Application Notes: Positive Control: Tonsil or lymph node (IHC).  
Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT) , (Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

Handling

is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images



SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified G-CSF Recombinant Rabbit Monoclonal Antibody (CSF3/3166R). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Tonsil stained with G-CSF Recombinant Rabbit Monoclonal Antibody (CSF3/3166R).