

Datasheet for ABIN6939235

Recombinant anti-Desmin antibody

Images




Overview

OVERVIEW	
Quantity:	100 μg
Target:	Desmin (DES)
Reactivity:	Human, Rat, Chicken, Hamster
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Desmin antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM), Immunostaining (ISt)
Product Details	
Immunogen:	Recombinant human full-length DES protein

Clone:	DES-2960R
Isotype:	IgG
Specificity:	Cytoskeletal intermediate filaments (IFs) constitute a diverse group of proteins that are
	expressed in a highly tissue-specific manner. IFs are constructed from two-chain -helical coiled-
	coil molecules arranged on an imperfect helical lattice, and have been widely used as markers
	for distinguishing individual cell types within a tissue and identifying the origins of metastatic
	tumors. Vimentin is an IF general marker of cells originating in the mesenchyme. Vimentin and
	Desmin, a related class III IF, are both expressed during skeletal muscle development. Desmin,
	a 469 amino acid protein found near the Z line in sarcomeres, is expressed more frequently in
	adult differentiated state tissues. Anti-desmin detects cells of normal smooth, skeletal, and


Product Details

Troduct Details	
	cardiac muscles.Antibody reacts with leiomyomas, leiomyosarcoma, rhabdomyomas,
	rhabdomyosarcoma, and perivascular cells of glomus tumors of the skin.
Purification:	Purified by Protein A/G
Target Details	
Target:	Desmin (DES)
Alternative Name:	DES (DES Products)
Molecular Weight:	52kDa
Gene ID:	1674
UniProt:	P17661
Application Details	
Application Notes:	Positive Control: SJRH30 cells. Uterus or Leiomyosarcoma.
	Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at
	RT)(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with
	1 mM EDTA, pH 9.0 for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for
	a specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
	is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified Desmin RabbitRecombinant Monoclonal Antibody (DES/2960R).Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human SkeletalMuscle stained with Desmin Rabbit RecombinantMonoclonal Antibody (DES/2960R).