

Datasheet for ABIN6939266
anti-Desmoglein 2 antibody[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	Desmoglein 2 (DSG2)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Desmoglein 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant human DSG2 protein
Clone:	6D8
Isotype:	IgG1 kappa
Specificity:	Recognizes a protein of 165 kDa, identified as Desmoglein-2 (DSG2). This monoclonal antibody recognizes the extracellular domain of human desmoglein-2. Desmoglein-2 is a member of the desmosomal cadherin family. Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. Desmocollins and desmogleins are the main desmosomal transmembrane proteins. Desmogleins consist of Dsg1, Dsg2, Dsg3, and Dsg4 isoforms. Within the desmosome, the extracellular domain of desmoglein is essential for calcium dependent heterophilic binding to the desmocollins, whereas the intracellular domain is essential for binding to the desmosomal plaque protein, plakoglobin. Human Desmoglein-2 is a type I transmembrane glycoprotein of 1117 amino acid (aa) residues with a 23 aa signal peptide and a 25 aa propeptide. It differs from other classic cadherins by having four instead of

Product Details

five cadherin repeat domains in its extracellular region, and a much larger cytoplasmic region containing five desmoglein repeat domains which share homology with the cadherin repeats. Instead of having the HAV adhesion motif found in type I cadherins, desmogleins have R/YAL as the adhesion motif on its amino-terminal cadherin repeat. The cytoplasmic tails of desmogleins interact with desmoplakins, plakoglobin and plakophilins. In turn, these proteins link the desmogleins with the intermediate filaments. Desmoglein-2 has been shown to be important in establishing cell-cell adhesion and function in epithelial cells. Desmoglein2 was originally identified in colon carcinoma and colon, and was named HDGC (human desmoglein colon).

Purification: Purified by Protein A/G

Target Details

Target: Desmoglein 2 (DSG2)

Alternative Name: DSG2 ([DSG2 Products](#))

Molecular Weight: 165kDa

Gene ID: 1829

UniProt: [Q14126](#)

Pathways: [Cell-Cell Junction Organization](#)

Application Details

Application Notes: Positive Control: A431, HepG2 or K562 cells. Heart or Colon.
Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (1-2 µg/mL), Western Blot (0.5-1.0 µg/mL) Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months

Images



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified DSG2 Mouse Monoclonal Antibody (6D8). Confirmation of Integrity and Purity of Antibody.