# antibodies -online.com





## Recombinant anti-ECM1 antibody

2 Images



#### Overview

Quantity:	100 μg
Target:	ECM1
Reactivity:	Human, Rat
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This ECM1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

#### **Product Details**

Immunogen:

iriiridi logeri.	Neodribinant namam fam Edwir protein
Clone:	ECM1-2889R
Isotype:	IgG
Specificity:	This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound
	Secretory Component. It does not react with the cell lines lacking secretory component. The
	antibody is useful for studying the distribution and level of both free and bound secretory
	component. Secretory component is differentially expressed in epithelium, and the antibody is a
	popular marker for identifying subpopulations of epithelial cells and epithelial differentiation.
	The Secretory component antibody is a useful research tool for studying mucosal immunity,
	inflammation, remodeling, differentiation and tumorigenesis, all processes associated with
	differential secretory component expression.


Recombinant human full-length ECM1 protein

### **Product Details** Purified by Protein A/G Purification: Target Details Target: ECM1 Alternative Name ECM1 (ECM1 Products) Molecular Weight: ~80kDa Gene ID: 1893 UniProt: Q16610 **Application Details** Positive Control: Stomach, Lung or breast tumor. **Application Notes:** Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT) (Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined. Restrictions: For Research Use only Handling Concentration: 200 μg/mL Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. 4 °C,-80 °C Storage: Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


24 months

**Expiry Date:** 



#### **SDS-PAGE**

Image 1. SDS-PAGE Analysis of Purified Secretory Component Rabbit Recombinant Monoclonal (ECM1/2889R). Confirmation of Purity and Integrity of Antibody.



#### **Immunohistochemistry**

**Image 2.** Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Secretory Component Rabbit Recombinant Monoclonal (ECM1/2889R).