

Datasheet for ABIN6939355  
**anti-Albumin antibody**



[Go to Product page](#)

3 Images

## Overview

Quantity:	100 µg
Target:	Albumin (ALB)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Albumin antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Coating (Coat), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant full-length human ALB protein
Clone:	ALB-2141
Isotype:	IgG1 kappa
Specificity:	<p>This MAb is absolutely specific to albumin and does not show any significant cross-reaction with other human proteins. Albumin is a soluble, monomeric protein, which comprises about one half of the blood serum protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Albumin is synthesized in the liver as preproalbumin, which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted form of albumin.</p>
Purification:	Purified by Protein A/G

## Target Details


Target:	Albumin (ALB)
Alternative Name:	ALB ( <a href="#">ALB Products</a> )
Molecular Weight:	66kDa
Gene ID:	213
UniProt:	<a href="#">P02768</a>
Pathways:	<a href="#">Lipid Metabolism</a>

## Application Details

Application Notes:	Positive Control: Human serum. Liver or Hepatocellular Carcinoma. Known Application: ELISA (For coating, order Ab without BSA),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

## Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months




**SDS-PAGE**

**Image 1.** SDS-PAGE Analysis Purified Albumin Mouse Monoclonal Antibody (ALB/2141). Confirmation of Purity and Integrity.

**Protein Array**

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using Albumin Mouse Monoclonal Antibody (ALB/2141). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



#### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Hepatocellular Carcinoma stained with Albumin Mouse Monoclonal Antibody (ALB/2141).