antibodies -online.com





anti-FCGR2A antibody



Image



Go to Product page

Overview

Quantity:	100 μg
Target:	FCGR2A
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FCGR2A antibody is un-conjugated
Application:	Immunofluorescence (IF), Flow Cytometry (FACS), Functional Studies (Func)

Product Details

Immunogen:	Recombinant human FCGR2A protein
Clone:	FCGR2A-479
Isotype:	IgG2b kappa
Purification:	Purified by Protein A/G

Target Details

Target:	FCGR2A
Alternative Name:	FCGR2A (FCGR2A Products)
Background:	This MAb binds most strongly to the IIa isoforms of CD32 (FcRII), a type 1 transmembrane
	glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and
	immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and
	at least 6 isoforms are generated via alternative mRNA splicing, i.e., Ila1, Ila2, Ilb1, Ilb2, Ilb3 and

Target Details

Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexes.
and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the
In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes
IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms.

Molecular Weight:	40kDa
Gene ID:	2212
UniProt:	P12318

Application Details


Application Notes:	Positive Control: K-562 cells. Lymph node, tonsil and placenta.
ADDIICATION NOTES	POSITIVE CONTOL K-SDZ CERS I VINDO NODE TONSILANO DIACENTA
, ipplication reces.	1 doitive doiti di. 11 doz dello. Eyrripii ridae, toridii aria piaderita.

Known Application: Functional Studies (Order Ab without BSA & Azide), Flow Cytometry (0.5-1 μ g/million cells), Immunofluorescence (0.5-1 μ g/mL), Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified CD32 Mouse Monoclonal Antibody (FCGR2A/479). Confirmation of Purity and Integrity of Antibody.