.-online.com antibodies

Datasheet for ABIN6939389 Recombinant anti-MLANA antibody

3 Images



Overview

Quantity:	100 µg
Target:	MLANA
Reactivity:	Human, Mouse, Rat, Dog
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This MLANA antibody is un-conjugated
Application:	Immunostaining (ISt), Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant full-length hMART-1 protein
Clone:	RMLANA-788
lsotype:	IgG1 kappa
Specificity:	This antibody recognizes a protein doublet of 20-22 kDa, identified as MART-1 (Melanoma
	Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte
	differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other
	melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1,
	MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that
	MART-1 is present in melanosomes and endoplasmic reticulum. This MAb labels melanomas
	and other tumors showing melanocytic differentiation. It is also a useful positive-marker for
	angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6939389 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Product Details		
	origin.	
Purification:	Purified by Protein A/G	
Target Details		
Target:	MLANA	
Alternative Name:	MLANA (MLANA Products)	

Molecular Weight:	20-22kDa (doublet)
Gene ID:	2315
UniProt:	Q16655
Application Details	
Application Notes:	Positive Control: SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues.
	Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at


Known Application: Immunonistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at
RT)(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Citrate
Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a
specific application should be determined.

Restrictions: For Research Use only

Handling


Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6939389 | 09/10/2023 | Copyright antibodies-online. All rights reserved.




Western Blotting

Image 1. Western Blot Analysis of COLO38 cell lysate using MART-1 Mouse Recombinant Monoclonal Antibody (rMLANA/788).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Melanoma stained with MART-1 Mouse Recombinant Monoclonal Antibody (rMLANA/788).



SDS-PAGE

Image 3. SDS-PAGE Analysis of Purified MART-1 MouseRecombinantMonoclonalAntibody(rMLANA/788).Confirmation of Purity and Integrity of Antibody.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6939389 | 09/10/2023 | Copyright antibodies-online. All rights reserved.