
antibodies .- online.com





anti-MLANA antibody





Go to Product page

Overview

Quantity:	100 μg
Target:	MLANA
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MLANA antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS), Immunostaining (ISt), Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:

Clone:	M2-9E3
Isotype:	IgG2b kappa
Specificity:	This antibody recognizes a protein doublet of 20-22 kDa, identified as MART-1 (Melanoma
	Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte
	differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other
	melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1,
	MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that
	MART-1 is present in melanosomes and endoplasmic reticulum. This MAb labels melanomas
	and other tumors showing melanocytic differentiation. It is also a useful positive-marker for
	angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal
	origin.


Recombinant hMART-1 protein

Product Details Purified by Protein A/G Purification: Target Details Target: **MLANA** MLANA (MLANA Products) Alternative Name: Molecular Weight: 20-22kDa (doublet) Gene ID: 2315 UniProt: Q16655 **Application Details** Positive Control: SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. **Application Notes:** Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (1-2 µg/mL), Western Blot (0.5-1 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined. Restrictions: For Research Use only Handling Concentration: 200 μg/mL 10 mM PBS with 0.05 % BSA & 0.05 % azide. Buffer: Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. Storage: 4 °C.-80 °C Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


24 months

Expiry Date:




Western Blotting

Image 1. Western Blot Analysis of COLO38 cell lysate using Melan-A MAb (M2-9E3).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Melanoma stained with Melan-A Monoclonal Antibody (M2-9E3).



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Melanoma stained with Melan-A Monoclonal Antibody (M2-9E3).

Please check the product details page for more images. Overall 4 images are available for ABIN6939392.