

Datasheet for ABIN6939432

Recombinant anti-AMACR antibody (AA 297-394)[Go to Product page](#)

4 Images

Overview

Quantity:	100 µg
Target:	AMACR
Binding Specificity:	AA 297-394
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This AMACR antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant human AMACR protein fragment (around aa 297-394) (exact sequence is proprietary)
Clone:	AMACR-2748R
Isotype:	IgG
Specificity:	This antibody recognizes a protein of 42 kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and -oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is expressed in cells of premalignant high-grade prostatic intraepithelial neoplasia (HGPIN) and prostate adenocarcinoma. The majority of the carcinoma cells show a distinct granular cytoplasmic staining reaction. AMACR is present at low or undetectable levels in glandular epithelial cells of

Product Details

normal prostate and benign prostatic hyperplasia. A spotty granular cytoplasmic staining is seen in a few cells of the benign glands. AMACR is expressed in normal liver (hepatocytes), kidney (tubular epithelial cells) and gall bladder (epithelial cells). Expression has also been found in lung (bronchial epithelial cells) and colon (colonic surface epithelium). AMACR expression can also be found in hepatocellular carcinoma and kidney carcinoma. Past studies have also shown that AMACR is expressed in various colon carcinomas (well, moderately and poorly differentiated) and over expressed in prostate carcinoma.

Purification: Purified by Protein A/G

Target Details

Target: AMACR

Alternative Name: AMACR ([AMACR Products](#))

Molecular Weight: 42kDa

Gene ID: 23600

UniProt: [Q9UHK6](#)

Pathways: [Monocarboxylic Acid Catabolic Process](#)

Application Details

Application Notes: Positive Control: HEK cells. Prostate Adenocarcinoma.
Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling


Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months

Images




Western Blotting

Image 1. Western Blot Analysis of human kidney tissue lysate using AMACR Recombinant Rabbit Monoclonal Antibody (AMACR/2748R).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with AMACR Recombinant Rabbit Monoclonal Antibody (AMACR/2748R).



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with AMACR Recombinant Rabbit Monoclonal Antibody (AMACR/2748R).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6939432.