



Datasheet for ABIN6939463

Recombinant anti-FTL antibody (AA 38-165)



[Go to Product page](#)

4 Images

Overview

Quantity:	100 µg
Target:	FTL
Binding Specificity:	AA 38-165
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This FTL antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant human FTL protein fragment (around aa 38-165) (exact sequence is proprietary)
Clone:	FTL-2338R
Isotype:	IgG
Purification:	Purified by Protein A/G

Target Details

Target:	FTL
Alternative Name:	FTL (FTL Products)
Background:	Mammalian ferritins consist of 24 subunits made up of 2 types of polypeptide chains, ferritin

Target Details

heavy chain and ferritin light chain. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe (II), whereas ferritin light chains promote the nucleation of ferrihydrite, enabling storage of Fe (III). Light chain ferritin is involved in cataracts by at least two mechanisms, hereditary hyperferritinemia cataract syndrome, in which light chain ferritin is overexpressed, and oxidative stress, an important factor in the development of ageing-related cataracts.

Molecular Weight: 19-25kDa

Gene ID: 2512

UniProt: [P02792](#)

Pathways: [Transition Metal Ion Homeostasis](#)

Application Details

Application Notes: Positive Control: HepG2, HeLa, HL-60 or 293T cells. Pancreas, Liver, Cerebellum or Testis.
Known Application: Western Blot (0.5-2 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1.0 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months




Western Blotting

Image 1. Western Blot Analysis of human kidney tissue lysate using Ferritin LC Rabbit Recombinant Monoclonal Antibody (FTL/2338R).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Ferritin LC Rabbit Recombinant Monoclonal Antibody (FTL/2338R).



SDS-PAGE

Image 3. SDS-PAGE Analysis Purified Ferritin LC Rabbit Recombinant Monoclonal Antibody (FTL/2338R). Confirmation of Purity and Integrity of Antibody.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6939463.