# antibodies - online.com







## anti-GP2 antibody (AA 35-179)





| $\sim$ |       |            |  |
|--------|-------|------------|--|
| ( )    | IV/Ar | $\sqrt{i}$ |  |

| Overview             |  |
|----------------------|--|
| Quantity:            | 100 μg   |
| Target:              | GP2  |
| Binding Specificity: | AA 35-179  |
| Reactivity:          | Human  |
| Host:                | Mouse  |
| Clonality:           | Monoclonal   |
| Conjugate:           | This GP2 antibody is un-conjugated   |
| Application:         | Immunohistochemistry (IHC), ELISA, Coating (Coat), Staining Methods (StM)                    |
| Product Details      |  |
| Immunogen:           | Recombinant fragment of human GP2 protein (around aa 35-179) (exact sequence is proprietary) |
| Clone:               | GP2-1712   |
| Isotype:             | IgG2b kappa  |
| No Cross-Reactivity: | Mouse (Murine)   |
| Purification:        | Purified by Protein A/G  |
| Target Details       |  |
| Target:              | GP2  |
| Alternative Name:    | GP2 (GP2 Products)   |
|                      |  |

### **Target Details**

Storage:

| Viral Protein   |  |
|---|--|
| GP2 (glycoprotein 2), also known as ZAP75, is a 537 amino acid secreted protein. It is an         |  |
| integral membrane protein that is secreted from intracellular zymogen granules and associates     |  |
| with the plasma membrane via glycosylphosphatidylinositol (GPI) linkage. GP2 is cleaved and       |  |
| then released into the pancreatic duct along with exocrine secretions. GP2 binds pathogens        |  |
| such as enterobacteria, thereby playing an important role in the innate immune response. The      |  |
| C-terminus of this protein is related to the C-terminus of the protein encoded by the neighboring |  |
| gene, uromodulin (UMOD). GP2 is also expressed on the apical plasma membrane of                   |  |
| specialized microfold (M) cells among enterocytes and serves as a transcytotic receptor for       |  |
| mucosal antigens. M cells are considered a promising target for oral vaccination against          |  |
| various infectious diseases.  |  |
| 59kDa   |  |
| 2813  |  |
| P55259  |  |
|   |  |
| Positive Control: PANC-1 cells. Pancreas.   |  |
| Known Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA),                |  |
| Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-      |  |
| fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min     |  |
| followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be    |  |
| determined.   |  |
| For Research Use only   |  |
|   |  |
| 200 μg/mL   |  |
| 10 mM PBS with 0.05 % BSA & 0.05 % azide.   |  |
|   |  |
| Sodium azide  |  |
| Sodium azide  This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which       |  |
| _   |  |


4 °C,-80 °C

#### Handling

| Storage Comment: | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody |
|------------------|---|
|                  | is stable for 24 months. Non-hazardous. No MSDS required.   |


Expiry Date: 24 months

#### **Images**




#### **Western Blotting**

**Image 1.** Western Blot Analysis of human Pancreas tissue lysate using GP2 Mouse Monoclonal Antibody (GP2/1712).



#### **Protein Array**

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using GP2 Mouse Monoclonal Antibody (GP2/1712) Z- and S- Score: The Zscore represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the



binding of that Monoclonal Antibody to protein X is equal to 29.

#### **Immunohistochemistry**

**Image 3.** Formalin-fixed, paraffin-embedded human Pancreas stained with GP2 Mouse Monoclonal Antibody (GP2/1712).

Please check the product details page for more images. Overall 4 images are available for ABIN6939542.