antibodies.com

# Datasheet for ABIN6939591 anti-GZMB antibody (AA 73-187)

5 Images



#### Overview

Quantity:	100 µg
Target:	GZMB
Binding Specificity:	AA 73-187
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This GZMB antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Staining Methods (StM)

### Product Details

Immunogen:	Recombinant fragment of human GZMB protein (around aa 73-187) (exact sequence is proprietary)
Clone:	GZMB-3014
Isotype:	IgG2b kappa
Purification:	Purified by Protein A/G

#### Target Details

Target:	GZMB
Alternative Name:	GZMB (GZMB Products)
Background:	Granzyme B is a member of the granule serine protease family stored specifically in NK cells or

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6939591 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

# Target Details

Molocular Woight:	20-2220
	shown to be an unfavorable prognostic indicator in Hodgkin's Disease.
	the identification of NK/T-cell lymphomas. High percentages of cytotoxic T-cells have been
	apoptosis by CTLs in the cell-mediated immune response. Granzyme B is useful as a marker in
	infection by intracellular pathogens. Granzyme B is crucial for the rapid induction of target cell
	cells bearing on their surface 'nonself' antigens, usually peptides or proteins resulting from
	recognize, bind, and lyse specific target cells. They are thought to protect their host by lysing
	cytotoxic T cells. Cytolytic T lymphocytes (CTL) and natural killer (NK) cells share the ability to

Molecular Weight:	29-32kDa
Gene ID:	3002
UniProt:	P10144
Pathways:	Apoptosis, Caspase Cascade in Apoptosis


## Application Details

Application Notes:	Positive Control: A431 cells. Human tonsil, spleen or Hodgkin's Lymphoma tissue (IHC).
	Known Application: Western Blot (1-2 $\mu$ g/mL), Immunohistochemistry (Formalin-fixed) (1-2 $\mu$
	g/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in
	10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal
	dilution for a specific application should be determined.
Restrictions:	For Research Use only

### Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6939591 | 09/10/2023 | Copyright antibodies-online. All rights reserved.




#### Western Blotting


**Image 1.** Western Blot Analysis of spleen tissue lysate using Granzyme B Mouse Monoclonal Antibody (GZMB/3014).

#### **Protein Array**

Image 2. Analysis of Protein Array containing >19,000 fulllength human proteins using Granzyme B Monospecific Mouse Monoclonal Antibody (GZMB/3014) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN6939591 | 09/10/2023 | Copyright antibodies-online. All rights reserved. Images



#### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Spleen stained with Granzyme B Monospecific Mouse Monoclonal Antibody (GZMB/3014).

Please check the product details page for more images. Overall 5 images are available for ABIN6939591.