

Datasheet for ABIN6939611

**Recombinant anti-DC-SIGN/CD209 antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	DC-SIGN/CD209 (CD209)
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This DC-SIGN/CD209 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunostaining (Ist), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant human CD209 protein fragment (exact sequence is proprietary)
Clone:	RC209-1781
Isotype:	IgG1 kappa
Specificity:	DC-SIGN is a transmembrane receptor that is expressed on the surface of dendritic cells and macrophages. It is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and

## Product Details

endogenous cells. The neck region is important for homo-oligomerization, which allows the receptor to bind multivalent ligands with high avidity.

Purification: Purified by Protein A/G

## Target Details

Target: DC-SIGN/CD209 (CD209)

Alternative Name: CD209 ([CD209 Products](#))

Molecular Weight: 45kDa

Gene ID: 30835

UniProt: [Q9NNX6](#)

## Application Details

Application Notes: Positive Control: THP-1 or HeLa cells (IF/FACS). Human liver or spleen lysate (WB). Human small intestine (IHC).

Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA Buffer, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months




SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified CD209 Mouse Monoclonal Antibody (rC209/1781). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CD209 Mouse Monoclonal Antibody (rC209/1781).



Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CD209 Mouse Monoclonal Antibody (rC209/1781).