

Datasheet for ABIN6939614

**anti-Neuregulin 1 antibody (AA 21-242)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	Neuregulin 1 (NRG1)
Binding Specificity:	AA 21-242
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Neuregulin 1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Coating (Coat)

## Product Details

Immunogen:	Recombinant fragment of human Neuregulin-1 (NRG1) protein (around aa 21-242) (exact sequence is proprietary)
Clone:	NRG1-2752
Isotype:	IgG
Purification:	Purified by Protein A/G

## Target Details

Target:	Neuregulin 1 (NRG1)
Alternative Name:	NRG1 ( <a href="#">NRG1 Products</a> )
Background:	Heregulin-1 is a membrane glycoprotein that mediates cell-cell signaling and plays a critical role

## Target Details

in the growth and development of multiple organ systems. An extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are expressed in a tissue-specific manner and differ significantly in their structure, and are classified as types I, II, III, IV, V and VI. Dysregulation of this gene has been linked to diseases such as cancer, schizophrenia, and bipolar disorder (BPD).

Molecular Weight:	26-71kDa
Gene ID:	3084
UniProt:	<a href="#">Q02297</a>
Pathways:	<a href="#">RTK Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Regulation of Muscle Cell Differentiation</a>


## Application Details

Application Notes:	Positive Control: A431, MCF-7 or T47D cells. Breast, Bladder or Thyroid Carcinoma. Known Application: ELISA (For coating, order Ab without BSA), Immunofluorescence (1-2 µg/mL)Optimal dilution for a specific application should be determined.
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months




SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752). Confirmation of Purity and Integrity of Antibody.

Protein Array

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



#### Immunofluorescence

**Image 3.** Confocal immunofluorescence of PFA-fixed MCF-7 cells using Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752) followed by Goat-anti Mouse CF488 (green) and Reddot is used to label the nuclei red.