

Datasheet for ABIN6939798

Recombinant anti-Insulin antibody**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Insulin (INS)
Reactivity:	Human, Pig, Mouse, Cow, Rabbit
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Insulin antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant human full-length Insulin protein (IRDN)
Clone:	IRDN-1980R
Isotype:	IgG
Specificity:	Recognizes a polypeptide which is identified as insulin, a 51-amino acid polypeptide composed of A and B chains connected through the C-peptide. Proinsulin, which has very little biological activity, is cleaved by proteases within its cell of origin into the insulin molecule and the C-terminal basic residue. Insulin enhances membrane transport of glucose, amino acids, and certain ions. It also promotes glycogen storage, formation of triglycerides, and synthesis of proteins and nucleic acids. Deficiency of insulin results in diabetes mellitus. The main storage site for insulin is the pancreatic islets. Antibodies to insulin are important as beta-cell and insulinoma marker.

Product Details

Purification: Purified by Protein A/G

Target Details

Target: Insulin (INS)

Alternative Name: INS ([INS Products](#))

Molecular Weight: 6kDa

Gene ID: 3630

UniProt: [P01308](#)

Pathways: [NF-kappaB Signaling](#), [RTK Signaling](#), [Positive Regulation of Peptide Hormone Secretion](#), [Peptide Hormone Metabolism](#), [Hormone Activity](#), [Carbohydrate Homeostasis](#), [ER-Nucleus Signaling](#), [Regulation of Carbohydrate Metabolic Process](#), [Feeding Behaviour](#), [Autophagy](#), [Negative Regulation of intrinsic apoptotic Signaling](#), [Brown Fat Cell Differentiation](#), [Positive Regulation of fat Cell Differentiation](#)

Application Details

Application Notes: Positive Control: MIA PaCa-2 cells. Pancreas.
Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months



SDS-PAGE

Image 1. SDS-PAGE Analysis of Purified Insulin Rabbit Recombinant Monoclonal Antibody (IRDN/1980R).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Pancreas stained with Insulin Rabbit Recombinant Monoclonal Antibody (IRDN/1980R).