antibodies .- online.com





anti-LGALS1/Galectin 1 antibody (AA 12-108)



Images



Go to Product page

_					
U	V	er	V	Ie	W

Alternative Name:

Quantity:	100 μg	
Target:	LGALS1/Galectin 1 (LGALS1)	
Binding Specificity:	AA 12-108	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This LGALS1/Galectin 1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Coating (Coat), Staining Methods (StM)	
Product Details		
Immunogen:	Recombinant fragment (around aa12-108) of human Galectin-1 protein (exact sequence is proprietary)	
Clone:	GAL1-1831	
Isotype:	IgG1 kappa	
Purification:	Purified by Protein A/G	
Target Details		
Target:	LGALS1/Galectin 1 (LGALS1)	

LGALS1 (LGALS1 Products)

Target Details

Background:

Galectin-1 is a member of the beta-galactoside-binding family and is a dimeric protein of 14kD participating in a variety of normal and pathological processes, including cancer progression. Galectin-1 can affect the proliferation of normal and malignant cells. Inhibition of cell growth is observed in a lactose-dependent manner as lower concentrations of the lectin stimulate cell proliferation. Galectin-1 may also be implicated in the induction of apoptosis of activated T cells through the binding of exogenous galectin-1 to CD45 Molecules present on the surface of lymphocytes. Galectin-1, reported to be present either at the surface of cancer cells or accumulated around these cells could act as an immunological shield to protect against a T cell immune response and provide an advantage for survival.

Molecular Weight:	14kDa
Gene ID:	3956
UniProt:	P09382

Application Details

۸nn	lication	Notos:
ADD	ucauon	ivotes.

Pathways:

Positive Control: HeLa, K562 or 293 cells. Prostate, Kidney, Placenta, Stomach, Skin, Spleen,

Brain or Heart.

Carbohydrate Homeostasis

Known Application: ELISA (Use Ab at 2-4 μ g/mL for coating) (Order Ab without BSA), Immunofluorescence (1-2 μ g/mL), Western Blot (1-2 μ g/mL), Immunohistochemistry (Formalinfixed) (0.5-1 μ g/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions:

For Research Use only

Handling

Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C


Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images




Western Blotting

Image 1. Western Blot Analysis of JEG-3 and K562 cell lysate using Galectin-1 Monospecific Mouse Monoclonal Antibody (GAL1/1831).



Protein Array

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using Galectin-1 Monospecific Mouse Monoclonal Antibody (GAL1/1831). Zand S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-lgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. Sscore therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Immunofluorescence

Image 3. Confocal immunofluorescence image of HeLa cells using Galectin-1 Monospecific Mouse Monoclonal Antibody (GAL1/1831). Green (CF488) and Reddot is used to label the nuclei Red.

Please check the product details page for more images. Overall 7 images are available for ABIN6939958.