

Datasheet for ABIN6939970 anti-LSP1 antibody (AA 174-278)

2 Images



Go to Product page

/ h	/er	1/1/	~ 1 A
1 11	/ 🗀 ।	1/16	→ \/\/
\cup	<i>/</i> C	V I V	\sim v v

Quantity:	100 μg
Target:	LSP1
Binding Specificity:	AA 174-278
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This LSP1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Coating (Coat)
Product Details	
Immunogen:	Recombinant fragment of human LSP1 protein (around aa174-278) (exact sequence is
Immunogen:	Recombinant fragment of human LSP1 protein (around aa174-278) (exact sequence is proprietary)
Immunogen: Clone:	
	proprietary)
Clone:	proprietary) LSP1-3042
Clone: Isotype:	proprietary) LSP1-3042 IgG1 kappa
Clone: Isotype: Purification:	proprietary) LSP1-3042 IgG1 kappa
Clone: Isotype: Purification: Target Details	proprietary) LSP1-3042 IgG1 kappa Purified by Protein A/G
Clone: Isotype: Purification: Target Details Target:	proprietary) LSP1-3042 IgG1 kappa Purified by Protein A/G LSP1

(including B and T lymphocytes, granulocytes and macrophages), phosphoprotein. However, mRNA splice variants that do not encode the lympho-specific protein are expressed from this gene in nonlymphoid cell lines as well (myocytes, stromal cells and fibroblasts), suggesting pp52 has a divergent role in signal transduction. The pp52 (LSP1) locus maps to human chromosome 11p15.5, which is implicated in tumor-related chromosomal translocations found in chronic lymphocytic leukemia. The pp52 promoter contains key elements that control transcriptional activity including an initiator specifying the unique 5' terminus of pp52 mRNA, tandem pairs of Ets and SP1 motifs, and a single C/EBP motif. LSP1 binds the cytoskeleton and has been implicated in affecting cytoskeletal remodeling in a variety of leukocyte functions, including cell motility and chemotaxis.

Molecular Weight:	52kDa	
Gene ID:	4046	
UniProt:	P33241	

specific application should be determined.


Positive Control: Human tonsil, colon, spleen or Hodgkin's lymphoma (IHC-P).

Known Application: ELISA (For coating, order antibody without BSA), Optimal dilution for a

Application Details


Application Notes:

	opcomo apprioation onodia de acterminea.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-80 °C	
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.	
Expiry Date:	24 months	



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified Monospecific Mouse Monoclonal Antibody to LSP1 (LSP1/3042). Confirmation of Purity and Integrity of Antibody.



Protein Array

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using LSP1 Mouse Monoclonal Antibody (LSP1/3042). Z- and S- Score: The Zscore represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Zscore, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.