antibodies

Datasheet for ABIN6939972 anti-TACSTD2 antibody (AA 31-274)

6 Images



Overview

Quantity:	100 µg
Target:	TACSTD2
Binding Specificity:	AA 31-274
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TACSTD2 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF), Staining Methods (StM)

Product Details

Immunogen:	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Clone:	TACSTD2-2152
Isotype:	IgG1 kappa
Purification:	Purified by Protein A/G


Target Details

Target:	TACSTD2
Alternative Name:	TACSTD2 (TACSTD2 Products)
Background:	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6939972 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details

	containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a
	phosphatidylinositol binding site and tyrosine phosphorylation sites near the C-terminus. It
	plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells,
	cornea and multi-stratified epithelia. It is also highly expressed in several types of tumors and is
	involved in regulating the growth of carcinoma cells.
Molecular Weight:	40kDa
Gene ID:	4070
UniProt:	P09758
Application Details	
Application Notes:	Positive Control: HT29 cells. Breast or Colon Carcinoma.
	Known Application: Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed)
	(1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections
	in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal
	dilution for a specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
	is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months




SDS-PAGE


Image 1. SDS-PAGE Analysis Purified TACSTD2 / TROP2MouseMonoclonalAntibody(TACSTD2/2152).Confirmation of Integrity and Purity of Antibody.

Protein Array

Image 2. Analysis of Protein Array containing >19,000 fulllength human proteins using TACSTD2 Mouse Monoclonal Antibody (TACSTD2/2152) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Zscore, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN6939972 | 09/10/2023 | Copyright antibodies-online. All rights reserved.



Immunofluorescence

Image 3. Confocal Immunofluorescence of MCF-7 cells TACSTD2 / TROP2 Mouse Monoclonal Antibody (TACSTD2/2152). labeled with CF488 (Green); Reddot is used to label the nuclei.

Please check the product details page for more images. Overall 6 images are available for ABIN6939972.