

Datasheet for ABIN6940053  
**anti-Ki-67 antibody (AA 2293-2478)**

## 8 Images

[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	Ki-67 (MKI67)
Binding Specificity:	AA 2293-2478
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Ki-67 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (IF), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant human Ki67 protein fragment (around aa 2293-2478) (exact sequence is proprietary)
Clone:	MKI67-2462
Isotype:	IgG2b kappa
Specificity:	<p>Ki-67 antigen is a nuclear, non-histone protein that is present in all stages of the cell cycle except G0. This characteristic makes Ki-67 an excellent marker for proliferating cells and is commonly used as one of the prognostic factors in cancer studies. A correlation has been demonstrated between Ki-67 index and the histo-pathological grade of neoplasms.</p> <p>Assessment of Ki-67 expression in renal and ureter tumors shows a correlation between tumor proliferation and disease progression, thus making it possible to differentiate high-risk patients.</p>

## Product Details

Ki-67 expression may also prove to be important for distinguishing between malignant and benign peripheral nerve sheath tumors. Ki-67 labeling index has been shown to be a prognostic marker in a number of neoplasms including grade II astrocytoma, oligodendroglioma, colon carcinoma, and breast carcinoma. In general, Ki-67 is a good marker of proliferating cell populations.

Purification: Purified by Protein A/G

## Target Details

Target: Ki-67 (MKI67)

Alternative Name: MKI67 ([MKI67 Products](#))

Molecular Weight: 345kDa and 395kDa

Gene ID: 4288

UniProt: [P46013](#)

Pathways: [Glycosaminoglycan Metabolic Process](#)

## Application Details

Application Notes: Positive Control: Any actively proliferating cells. Skin, Tonsil or Lymph Node.  
Known Application: Western Blot (1-2 µg/mL), Flow Cytometry (1-2 µg/million cells), Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-80 °C

Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months

Images





Western Blotting

**Image 1.** Western Blot Analysis of Raji and Ramos cell lysate using Ki67-Monospecific Mouse Monoclonal Antibody (MKI67/2462).

Protein Array

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using Ki67-Monospecific Mouse Monoclonal Antibody (MKI67/2462). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.





### Flow Cytometry

**Image 3.** Flow Cytometric Analysis of HeLa cells. Ki67-Monospecific Mouse Monoclonal Antibody (MKI67/2462) followed by goat anti-Mouse IgG-CF488 (blue); Isotype Control (red).

Please check the [product details page](#) for more images. Overall 8 images are available for ABIN6940053.