antibodies.com

Datasheet for ABIN6940060 anti-MME antibody (AA 297-483)

5 Images



Overview

Quantity:	100 µg
Target:	MME
Binding Specificity:	AA 297-483
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MME antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Staining Methods (StM), Coating (Coat)

Product Details

Immunogen:	Recombinant human CD10 protein fragment (aa297-483) (exact sequence is proprietary)
Clone:	MME-1892
Isotype:	lgG2c kappa
Specificity:	Recognizes a 100 kDa glycoprotein, identified as CD10, also known as Common Acute
	Lymphocytic Leukemia Antigen (CALLA). It is a cell surface enzyme with neutral
	metalloendopeptidase activity, which inactivates a variety of biologically active peptides. CD10
	is expressed on the cells of lymphoblastic, Burkitt's, and follicular germinal center lymphomas,
	and on cells from patients with chronic myelocytic leukemia (CML). It is also expressed on the
	surface of normal early lymphoid progenitor cells, immature B cells within adult bone marrow
	and germinal center B cells within lymphoid tissue. CD10 is also present on breast

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6940060 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Product Details	
	myoepithelial cells, bile canaliculi, fibroblasts, with especially high expression on the brush border of kidney and gut epithelial cells.
Purification:	Purified by Protein A/G
Target Details	
Target:	MME
Alternative Name:	MME (MME Products)

by Hormones, Smooth Muscle Cell Migration

RTK Signaling, Peptide Hormone Metabolism, Regulation of Systemic Arterial Blood Pressure

Application	Details
	2 0 000

Target Type:

Gene ID:

UniProt:

Pathways:

Molecular Weight:

Chemical

100kDa

P08473

4311

Application Notes:	Positive Control: Tonsil, small intestine or kidney (IHC).	
	Known Application: ELISA (Use Ab at 2-4 μ g/mL for coating) (Order Ab without BSA), Western	
	Blot (1-2 μ g/mL),Immunohistochemistry (Formalin-fixed) (1-2 μ g/mL for 30 minutes at	
	RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM	
	EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a	
	specific application should be determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 µg/mL	
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-80 °C	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6940060 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Handling


Storage Comment:

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date:

24 months

Images



Western Blotting

Image 1. Western Blot Analysis of kidney tissue lysate using CD10 Mouse Monoclonal Antibody (MME/1892).


. .

Protein Array

104 Strength of Signal (Z score) 78 TRIM24 1.73 0.34 52 1.39 0.17 26 0 10 20 30 40 50 Signal Rank (Top 50)

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using CD10 Mouse Monoclonal Antibody (MME/1892) Z- and S- Score: The Zscore represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Zscore, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Images



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Kidney stained with CD10 Mouse Monoclonal Antibody (MME/1892).

Please check the product details page for more images. Overall 5 images are available for ABIN6940060.