

Datasheet for ABIN6940181

anti-NEFH antibody

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	NEFH
Reactivity:	Human, Rat, Mouse, Chicken, Pig
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This NEFH antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Staining Methods (StM), Coating (Coat)

Product Details


Immunogen:	Human recombinant neurofilament protein
Clone:	SPM563
Isotype:	IgG1 kappa
Specificity:	This MAb reacts with a 200 kDa protein, identified as heavy sub-unit of neurofilaments (NF-H).
	Neurofilaments make up the main structural elements of axons and dendrites and are found in
	neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three
	major subunits with molecular weights of 68 kDa (NF-L), 160 kDa (NF-M) and 200 kDa (NF-H).
	Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors.
	Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas
	stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as
	well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine
	carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.

Product Details Purification: Purified by Protein A/G Target Details Target: **NEFH NEFH (NEFH Products)** Alternative Name: Molecular Weight: 200kDa Gene ID: 4744 UniProt: P12036 **Application Details** Positive Control: Brain, Neuroblastoma. **Application Notes:** Known Application: ELISA (For coating, order Ab without BSA), Flow Cytometry (0.5-1 µg/million cells), Western Blot (0.5-1 μg/mL), Immunohistochemistry (Frozen & formalin-fixed) (0.5-1 μ g/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined. Restrictions: For Research Use only Handling Concentration: 200 μg/mL 10 mM PBS with 0.05 % BSA & 0.05 % azide. Buffer: Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. Storage: 4 °C.-80 °C Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


24 months

Expiry Date:



Western Blotting

Image 1. Western Blot Analysis of human brain tissue lysate using Neurofilament MAb (SPM563).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded humanCerebellum stained with Neurofilament MonoclonalAntibody (SPM563).