

Datasheet for ABIN6940210  
**anti-NKX3-1 antibody (AA 92-224)**[Go to Product page](#)

## 4 Images

## Overview

Quantity:	100 µg
Target:	NKX3-1
Binding Specificity:	AA 92-224
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This NKX3-1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Coating (Coat), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant fragment (around aa 92-224) of human NKX3.1 protein (exact sequence is proprietary)
Clone:	NKX3-1-2836
Isotype:	IgG1 kappa
Purification:	Purified by Protein A/G

## Target Details

Target:	NKX3-1
Alternative Name:	NKX3-1 ( <a href="#">NKX3-1 Products</a> )
Background:	NKX3.1 is a prostate specific gene encoding a transcription factor that plays an important role

## Target Details

in normal prostate development and carcinogenesis. It is a prostatic tumor suppressor gene located on chromosome 8p21.2, which frequently undergoes a loss of heterozygosity. NKX3.1 expression is highly restricted in prostate epithelial cells and therefore can be used as a diagnostic biomarker for prostate cancer and other metastatic lesions of prostatic origin. Furthermore, NKX3.1 shows better sensitivity than Prostate Specific Antigen (PSA) for identifying metastatic prostatic adenocarcinoma. This suggests that immunohistochemical staining of NKX3.1, along with other prostate-restricted markers, may be valuable for the definitive determination of prostatic origin in poorly differentiated metastatic carcinomas.

Molecular Weight: 35kDa

Gene ID: 4824

UniProt: [Q99801](#)

Pathways: [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Positive Regulation of Endopeptidase Activity](#), [Positive Regulation of Response to DNA Damage Stimulus](#)

## Application Details

Application Notes: Positive Control: Highly expressed in the prostate and at a lower level in the testis.  
Known Application: ELISA (For coating, order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT), (Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months




SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified NKX3.1-Monospecific Mouse Monoclonal Antibody (NKX3.1/2836). Confirmation of Purity and Integrity of Antibody.



Protein Array

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using NKX3.1-Monospecific Mouse Monoclonal Antibody (NKX3.1/2836). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



#### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human prostate stained with NKX3.1-Monospecific Mouse Monoclonal Antibody (NKX3.1/2836).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6940210.