# antibodies - online.com







# Recombinant anti-YBX1 antibody

**Images** 



#### Overview

| Quantity:      | 100 μg   |
|----------------|--|
| Target:        | YBX1   |
| Reactivity:    | Human  |
| Host:          | Mouse  |
| Antibody Type: | Recombinant Antibody                               |
| Clonality:     | Monoclonal   |
| Conjugate:     | This YBX1 antibody is un-conjugated                |
| Application:   | Immunohistochemistry (IHC), Staining Methods (StM) |

## **Product Details**


| Immunogen:    | Recombinant human full-length YBX1 protein |
|---------------|--|
| Clone:        | RYBX1-2430                                 |
| Isotype:      | IgG1 kappa                                 |
| Purification: | Purified by Protein A/G                    |

# Target Details

| Target:           | YBX1  |
|-------------------|---|
| Alternative Name: | YBX1 (YBX1 Products)  |
| Background:       | The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. Y-box binding protein-1 (YBX1) |


| Target Details      |  |
|---------------------|--|
|                     | is the prototypic member of the cold shock protein family that fulfills numerous cellular functions. In the nucleus, YBX1 protein orchestrates transcription of proliferation-related genes, whereas in the cytoplasm it associates with mRNA and directs translation. In human tumor entities, such as breast, lung and prostate cancer, cellular YBX1 expression indicates poor clinical outcome, suggesting that YBX1 is an attractive marker to predict patients' prognosis. |
| Molecular Weight:   | 36kDa  |
| Gene ID:            | 4904   |
| UniProt:            | P67809   |
| Pathways:           | Regulation of Muscle Cell Differentiation  |
| Application Details |  |
| Application Notes:  | Positive Control: MCF-7, HEK293 or PC12 cells. Prostate, Breast or Lung Carcinoma.  Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Concentration:      | 200 μg/mL  |
| Buffer:             | 10 mM PBS with 0.05 % BSA & 0.05 % azide   |

| Concentration:     | 200 μg/mL   |
|--------------------|---|
| Buffer:            | 10 mM PBS with 0.05 % BSA & 0.05 % azide.   |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                                      |
| Storage:           | 4 °C,-80 °C   |
| Storage Comment:   | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required. |
| Expiry Date:       | 24 months   |



### **SDS-PAGE**

**Image 1.** SDS-PAGE Analysis Purified YBX1 Mouse Recombinant Monoclonal Antibody (rYBX1/2430). Confirmation of Purity and Integrity of Antibody.



#### **Immunohistochemistry**

**Image 2.** Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with YBX1 Mouse Recombinant Monoclonal Antibody (rYBX1/2430).