antibodies -online.com





anti-PODXL antibody (Extracellular Domain)





0 -	4.4	Prod	400		
1-()	10	Pron	ויאוו	nac	\Box
\circ	LO.	1 100	uot	Duc	\sim

Overview		
Quantity:	100 μg	
Target:	PODXL	
Binding Specificity:	Extracellular Domain	
Reactivity:	Human, Rat, Rabbit	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This PODXL antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Flow Cytometry (FACS),	
	Immunofluorescence (IF), Immunoprecipitation (IP), Staining Methods (StM)	
Product Details		
Product Details Immunogen:	A recombinant protein fragment containing the intracellular, transmembrane, and part of the	
	A recombinant protein fragment containing the intracellular, transmembrane, and part of the extracellular domain of human podocalyxin	
Immunogen:	extracellular domain of human podocalyxin	
Immunogen: Clone:	extracellular domain of human podocalyxin 3D3	
Immunogen: Clone: Isotype:	extracellular domain of human podocalyxin 3D3 IgG1 kappa	
Immunogen: Clone: Isotype: Purification:	extracellular domain of human podocalyxin 3D3 IgG1 kappa	

Target Details


rarget betails			
Background:	Podocalyxin is a member of the CD34 transmembrane sialomucin family. It is over-expressed on the podocyte foot projections and plays essential roles in kidney development and homeostasis, blood filtration and urine formation. It is also expressed on vascular endothelia, hematopoietic progenitors and a subset of neurons. Overexpression of podocalyxin may be linked to more aggressive tumor behavior. Podocalyxin antibody can identify podocytes in the urine (podocyturia) that may indicate glomerular disease, pre-eclampsia, and other kidney pathology.		
Molecular Weight:	165-170kDa		
Gene ID:	5420		
UniProt:	000592		
Pathways:	Tube Formation		
Application Details			
Application Notes:	Positive Control: HeLa, Raji, Jurkat cells, Angiosarcoma, Breast, Prostate, Liver, Pancreatic, Kidney. Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (1-2 µg/mL) Immunoprecipitation (1-2 µg/500 µg protein), Western Blot (0.5-1 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.		
Restrictions:	For Research Use only		
Handling			
Concentration:	200 μg/mL		
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	4 °C,-80 °C		
Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C			

is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date:


24 months

Images




Western Blotting

Image 1. Western Blot Analysis of HeLa cell lysate using Podocalyxin Mouse Monoclonal Antibody (3D3).



Immunofluorescence

Image 2. Confocal Immunofluorescence of HeLa cells using Podocalyxin Mouse Monoclonal Antibody (3D3) labeled with CF488 (Green); Reddot is used to label the nuclei.



Flow Cytometry

Image 3. Flow Cytometry of NCCIT cells using Podocalyxin Mouse Monoclonal Antibody (3D3).

Please check the product details page for more images. Overall 6 images are available for ABIN6940326.