antibodies -online.com





Recombinant anti-ANO1 antibody (AA 2-101)

ANO1

TMEM16A (ANO1 Products)

2 Images

Overview

Target:

Alternative Name:



Go to Product page


Quantity:	100 μg
Target:	ANO1
Binding Specificity:	AA 2-101
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This ANO1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)
Product Details	
Immunogen:	A recombinant human DOG-1 protein fragment (around aa 2-101) (exact sequence is proprietary)
Clone:	DG1-2564R
Isotype:	IgG
Purification:	Purified by Protein A/G
Target Details	

Target Details

Expiry Date:


rarget Details	
Background:	Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST's), c-kit
	signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft
	tissue tumors, which, due to appearance, may be difficult to diagnose. Immunoreactivity for
	DOG-1 has been reported in 97.8 percent of scorable GIST's, including all c-kit negative GIST's
	Overexpression of DOG-1 has been suggested to aid in the identification of GISTs, including
	Platelet-Derived Growth Factor Receptor Alpha mutants that fail to express c-kit antigen. The
	overall sensitivity of DOG1 and c-kit in GIST's is nearly identical: 94.4 % vs. 94.7 %.
Molecular Weight:	~114kDa
Gene ID:	55107
Application Details	
Application Notes:	Positive Control: Gastrointestinal Stromal Tumor (GIST) or testicular germ cell tumor.
	Melanocytes in the basal layer of the epidermis and mast cells in the dermis of normal skin.
	Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at
	RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate
	buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a
	specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
	is stable for 24 months. Non-hazardous. No MSDS required.

24 months



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified DOG-1 RabbitRecombinant Monoclonal Antibody (DG1/2564R).Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human GIST stained with DOG-1 Rabbit Recombinant Monoclonal Antibody (DG1/2564R).