

Datasheet for ABIN6940384

anti-PPP3R1 antibody[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	PPP3R1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PPP3R1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant full-length Calcineurin B protein
Clone:	CALNB-2342
Isotype:	IgG1 kappa
Purification:	Purified by Protein A/G

Target Details

Target:	PPP3R1
Alternative Name:	PPP3R1 (PPP3R1 Products)
Background:	Calcineurin is an enzyme that dephosphorylates serine and threonine residues in proteins. It is a heterodimer of a 59 kDa catalytic A subunit and a 19 kDa regulatory B subunit that is activated by the binding of calcium ions and calmodulin. Calcineurin is expressed in many tissues, but its levels are highest in the brain, where it may play a role in learning and memory. It has many

Target Details

substrates, including NFAT, a transcription factor that is activated by dephosphorylation. Complexes of the immuno-suppressants cyclosporine and FK506 with immunophilin proteins such as cyclophilin and FKBP12 are potent and specific inhibitors of Calcineurin activity. Alterations in Calcineurin activity are suspected to play a role in cardiac hypertrophy and graft versus host disease in organ transplantation.


Molecular Weight:	19kDa
Gene ID:	5534
UniProt:	P63098
Pathways:	RTK Signaling , WNT Signaling , Fc-epsilon Receptor Signaling Pathway , Protein targeting to Nucleus , VEGF Signaling , BCR Signaling

Application Details

Application Notes:	Positive Control: HeLa, MCF-7 or A431 cells. Prostate, Thyroid or Pancreas. Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



Immunohistochemistry

Image 1. Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Calceurin B Mouse Monoclonal Antibody (CALNB/2342).