

Datasheet for ABIN6940442

anti-CD45 antibody**4** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CD45 (PTPRC)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD45 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Western Blotting (WB), Immunofluorescence (IF), Staining Methods (StM)

Product Details

Immunogen:	Human B-lymphocytes
Clone:	PTPRC-1132
Isotype:	IgG1 kappa
No Cross-Reactivity:	Monkey, Rat (Rattus)
Purification:	Purified by Protein A/G

Target Details

Target:	CD45 (PTPRC)
Alternative Name:	PTPRC (PTPRC Products)
Background:	CD45R, also designated CD45 and PTPRC, is identified as a transmembrane glycoprotein,

Target Details

broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.

Molecular Weight: 180-220kDa

Gene ID: 5788

UniProt: [P08575](#)

Pathways: [TCR Signaling](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [CXCR4-mediated Signaling Events](#), [BCR Signaling](#)

Application Details

Application Notes: Positive Control: Ramos, U-698 or GA-10 cells. Spleen and Tonsil.
Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (0.5-1 µg/mL), Western Blot (0.5-1.0 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1.0 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

Handling

is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images




Western Blotting

Image 1. Western Blot Analysis of human Spleen tissue lysates using CD45RB Monoclonal Antibody (PTPRC/1132).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Lymphoma stained with CD45RB Monoclonal Antibody (PTPRC/1132).



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Tonsil stained with CD45RB Monoclonal Antibody (PTPRC/1132).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6940442.