

Datasheet for ABIN6940477

anti-RAD51 antibody (AA 1-134)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	RAD51
Binding Specificity:	AA 1-134
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This RAD51 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Coating (Coat)

Product Details

Immunogen:	Recombinant fragment of human RAD51 protein (around aa 1-134) (exact sequence is proprietary)
Clone:	RAD51-2753
Isotype:	IgG
Purification:	Purified by Protein A/G

Target Details

Target:	RAD51
Alternative Name:	RAD51 (RAD51 Products)
Background:	RAD51 is one of the key factors of DNA repair by homologous recombination and has been

Target Details

shown to have anti-apoptotic activity in tumor cells. RAD51 protein interacts with a variety of tumor suppressor proteins including p53, BRCA1 and BRCA2. Elevated expression of RAD51 enhances radio-resistance of human tumor cells. Overexpression of RAD51 protein in tumor cells renders them resistant against cytotoxic drugs like Cisplatin. RAD51 interacts with BRCA1 and BRCA2 to influence subcellular localization and cellular response to DNA damage. BRCA2 inactivation may be a key event leading to genomic instability and tumorigenesis from deregulation of RAD51. High-level expression of RAD51 has been observed in a variety of human malignancies. RAD51 overexpression correlates with histological grading of the tumor in invasive ductal mammary carcinoma.


Molecular Weight:	37kDa
Gene ID:	5888
UniProt:	Q06609
Pathways:	DNA Damage Repair

Application Details

Application Notes:	Positive Control: HeLa or MCF-7 cells. Colon or breast Carcinoma. Known Application: ELISA (For coating, order Ab without BSA), Western Blot (1-2 µg/mL) Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



Western Blotting

Image 1. Western Blot Analysis of K562, HeLa and MCF-7 cell lysates using RAD51 Mouse Monoclonal Antibody (RAD51/2753).

Protein Array

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using RAD51 Mouse Monoclonal Antibody (RAD51/2753) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.