



Datasheet for ABIN6940573  
**anti-VISTA antibody**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	VISTA
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This VISTA antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Coating (Coat), Staining Methods (StM)

Product Details

Immunogen:	Recombinant full-length human VISTA protein
Clone:	VISTA-2865
Isotype:	IgG2a kappa
Purification:	Purified by Protein A/G

Target Details

Target:	VISTA
Alternative Name:	VSIR ( <a href="#">VISTA Products</a> )
Background:	VISTA / Gi24 is a transmembrane protein expressed in bone, on embryonic stem cells (ESCs), and on tumor cell surfaces. On ESC s, Gi24 appears to positively interact with BMP-4, potentiating BMP signaling and the transition from an undifferentiated to a differentiated state. On tumor cells, Gi24 both promotes MT1-MMP expression and activity and serves as a

## Target Details

---

substrate for MT1-MMP. This increases the potential for cell motility. Mature human Gi24 contains a 162aa extracellular region with one V-type Ig-like domain and a 96aa cytoplasmic domain. Human Gi24 undergoes proteolytic cleavage by MT1-MMP, generating a soluble 30 kDa extracellular fragment plus a 25-30 kDa membrane-bound fragment. VISTA is a negative checkpoint regulator and is expressed on myeloid cells, T-cells and human TILs (tumor infiltrating lymphocytes) on MDSCs (myeloid-derived suppressor cells) in the TME (tumor microenvironment). It is very likely both a ligand and receptor and is a promising target for cancer immunotherapy.

Molecular Weight:	34kDa
Gene ID:	64115
UniProt:	<a href="#">Q9H7M9</a>
Pathways:	<a href="#">Cancer Immune Checkpoints</a>

## Application Details


---

Application Notes:	Positive Control: Human spleen or placenta tissue (IHC). Known Application: ELISA (For coating, order antibody without BSA), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

## Handling


---

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



### SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified Monospecific Mouse Monoclonal Antibody to VISTA (VISTA/2865). Confirmation of Integrity and Purity of Antibody.



### Protein Array

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using Monospecific Mouse Monoclonal Antibody to VISTA (VISTA/2865). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.