.-online.com antibodies

Datasheet for ABIN6940576 anti-SHBG antibody

2 Images



Overview

Quantity:	100 µg
Target:	SHBG
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SHBG antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Coating (Coat), Staining Methods (StM)

Product Details

Immunogen:	Recombinant full-length human SHBG protein
Clone:	SHBG-245
Isotype:	IgG1 kappa
Specificity:	Recognizes a protein of 45 kDa, identified as SHBG. It functions as an androgen transport
	protein, but may also be involved in receptor-mediated processes. Each dimer binds one
	molecule of steroid. It is specific for 5-alpha-dihydrotestosterone, testosterone, and 17-beta-
	estradiol. SHBG regulates the plasma metabolic clearance rate of steroid hormones by
	controlling their plasma concentration. In testis, it is synthesized by the Sertoli cells, secreted
	into the lumen of the seminiferous tubule and transported to the epididymis.
Purification:	Purified by Protein A/G

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6940576 | 09/10/2023 | Copyright antibodies-online. All rights reserved.


Target Details

Target:	SHBG
Alternative Name:	SHBG (SHBG Products)
Molecular Weight:	45kDa
Gene ID:	6462
UniProt:	P04278
Pathways:	Hormone Transport

Application Details


Positive Control: Liver, Placenta or Testis.
Known Application: ELISA (For coating, order Ab without BSA), ,Immunohistochemistry
(Formalin-fixed) (1-2 μ g/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires
boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by
cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
For Research Use only
200 µg/mL
10 mM PBS with 0.05 % BSA & 0.05 % azide.
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
should be handled by trained staff only.
4 °C,-80 °C
Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
is stable for 24 months. Non-hazardous. No MSDS required.
24 months

Images



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified SHBG Mouse Monoclonal Antibody (SHBG/245). Confirmation of Integrity and Purity of Antibody.



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Testicular Carcinoma stained with SHBG Mouse Monoclonal Antibody (SHBG/245).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6940576 | 09/10/2023 | Copyright antibodies-online. All rights reserved.