

Datasheet for ABIN6940628

**Recombinant anti-SOX10 antibody (AA 115-269)**[Go to Product page](#)

## 4 Images

## Overview

Quantity:	100 µg
Target:	SOX10
Binding Specificity:	AA 115-269
Reactivity:	Human, Mouse
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This SOX10 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	A recombinant fragment (around aa115-269) of human SOX10 protein (exact sequence is proprietary)
Clone:	SOX10-2311R
Isotype:	IgG
Purification:	Purified by Protein A/G

## Target Details

Target:	SOX10
Alternative Name:	SOX10 ( <a href="#">SOX10 Products</a> )

## Target Details

Background:	<p>SOX genes comprise a family of genes that are related to the mammalian sex-determining gene SRY. These genes similarly contain sequences that encode for the HMG-box domain, which is responsible for the sequence-specific DNA-binding activity. SOX-10 is a sensitive marker of melanoma, including conventional, spindled, and desmoplastic subtypes. It is expressed by metastatic melanomas and nodal capsular nevus in sentinel lymph nodes, but not by other lymph node components such as dendritic cells, which usually express S100 protein.</p> <p>Commonly used melanoma markers, such as anti-HMB-45 and anti-Melan-A, are poorly expressed in desmoplastic melanomas while SOX-10 is moderately to strongly expressed in desmoplastic melanomas. SOX-10 is considered as a very reliable marker for recognizing residual desmoplastic melanomas. In normal tissues, it is expressed in Schwann cells, melanocytes, and myoepithelial cells of salivary, bronchial and mammary glands. SOX-10 expression is also observed in mast cells.</p>
Molecular Weight:	49-58kDa
Gene ID:	6663
UniProt:	<a href="#">P56693</a>
Pathways:	<a href="#">Chromatin Binding</a>

## Application Details

Application Notes:	<p>Positive Control: HepG2 cells. Melanomas, breast carcinomas, gliomas.</p> <p>Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.</p>
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling


Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C

Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images



Western Blotting


**Image 1.** Western Blot Analysis of COLO-38 cell lysate using SOX10-Monospecific Recombinant Rabbit Monoclonal Antibody (SOX10/2311R).



Protein Array

**Image 2.** Analysis of Protein Array containing more than 19,000 full-length human proteins using SOX10-Monospecific Rabbit Monoclonal Antibody (SOX10/2311R)  
Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the

S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Melanoma stained with SOX10-Monospecific Recombinant Rabbit Monoclonal Antibody (SOX10/2311R).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6940628.