antibodies.com

# Datasheet for ABIN6940686 anti-TFF1 antibody (AA 57-84)

3 Images



## Overview

Quantity:	100 µg
Target:	TFF1
Binding Specificity:	AA 57-84
Reactivity:	Human, Cynomolgus
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TFF1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	A synthetic peptide (around aa 57-84) of human pS2/TFF1 polypeptide (exact sequence is proprietary)
Clone:	GE2-same as R47-94
lsotype:	IgG1 kappa
Specificity:	It recognizes a polypeptide of 6.5 kDa, identified as pS2 estrogen-regulated protein. Its epitope
	is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are
	protease resistant molecules secreted throughout the gut that play a role in mucosal healing.
	These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain.
	pS2 is known to form dimers and this dimerization is thought to play a role in its protective and
	healing properties. About 60 % of breast carcinomas are positive for pS2. Staining is
	cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6940686 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

FIGUUCI Details	
	normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in
	ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily
	expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen
	dependent tumors that displays an increased likelihood of response to endocrine therapy.
Purification:	Purified by Protein A/G
Target Details	
Target:	TFF1
Alternative Name:	TFF1 (TFF1 Products)
Molecular Weight:	6.5kDa
Gene ID:	7031
UniProt:	P04155
Pathways:	EGFR Signaling Pathway
Application Details	
Application Notes:	Positive Control: Breast or Ovarian carcinoma.
	Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1.0 $\mu$ g/mL for 30 minutes at
	RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate
	Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a
	specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6940686 | 09/10/2023 | Copyright antibodies-online. All rights reserved.


## Handling

is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date:


#### 24 months

## Images



#### SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified pS2 Mouse Monoclonal Antibody (GE2). Confirmation of Purity and Integrity of Antibody.



### Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with pS2 Mouse Monoclonal Antibody (GE2).

## Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with pS2 Mouse Monoclonal Antibody (GE2).