

Datasheet for ABIN6940692

Recombinant anti-TFF1 antibody (AA 57-84)[Go to Product page](#)**2** Images

Overview

Quantity:	100 µg
Target:	TFF1
Binding Specificity:	AA 57-84
Reactivity:	Human, Cynomolgus
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This TFF1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human pS2.
Clone:	TFF1-2969R
Isotype:	IgG
Specificity:	<p>It recognizes a polypeptide of 6.5 kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing.</p> <p>These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60 % of breast carcinomas are positive for pS2. Staining is</p>

Product Details

cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.

Purification: Purified by Protein A/G

Target Details

Target: TFF1

Alternative Name: TFF1 ([TFF1 Products](#))

Molecular Weight: 6.5kDa

Gene ID: 7031

UniProt: [P04155](#)

Pathways: [EGFR Signaling Pathway](#)

Application Details

Application Notes: Positive Control: Breast or Ovarian carcinoma.
Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1.0 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-80 °C

Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images



SDS-PAGE

Image 1. SDS-PAGE Analysis Purified pS2 Rabbit Recombinant Monoclonal Antibody (TFF1/2969R). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human Stomach Carcinoma stained with pS2 Rabbit Recombinant Monoclonal Antibody (TFF1/2969R).