

Datasheet for ABIN6940716  
**anti-TGFA antibody (C-Term)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µg
Target:	TGFA
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Mouse, Cow
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TGFA antibody is un-conjugated
Application:	Immunofluorescence (IF), Flow Cytometry (FACS)

## Product Details

Immunogen:	A 17-amino acid synthetic peptide from the carboxyl-terminus of rat TGFα.
Clone:	SPM357
Isotype:	IgG1 kappa
Specificity:	This antibody reacts with the C-terminus of TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with Ab-1 is completely blocked by the peptide used for raising this antibody. TGF (aa50) is a growth factor with 33 % homology to EGF, binds to EGFR, activates tyrosine phosphorylation of the receptor, and stimulates cell proliferation. It plays a role in tumor initiation by inducing the reversible transformed phenotype.
Purification:	Purified by Protein A/G

## Target Details


Target:	TGFA
Alternative Name:	TGFA ( <a href="#">TGFA Products</a> )
Molecular Weight:	~6kDa
Gene ID:	7039
UniProt:	<a href="#">P01135</a>
Pathways:	<a href="#">NF-kappaB Signaling</a> , <a href="#">RTK Signaling</a> , <a href="#">EGFR Signaling Pathway</a>

## Application Details

Application Notes:	Positive Control: Jurkat or Ramos cells. Testicular Carcinoma. Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (1-2 µg/mL), Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

## Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



#### Immunohistochemistry

**Image 1.** Formalin-fixed, paraffin-embedded human Testicular Carcinoma stained with TGF alpha Mouse Monoclonal Antibody (SPM357)