# antibodies .- online.com





## anti-Tyrosinase-Related Protein 1 antibody

2 Images



#### Overview

Quantity:	100 μg
Target:	Tyrosinase-Related Protein 1 (TYRP1)
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tyrosinase-Related Protein 1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

#### **Product Details**

Immunogen:

Isotype:	IgG
Specificity:	It reacts with a 75 kDa melanocyte-specific gene product, identified as Tyrosinase-related
	protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal
	membranes of melanoma, normal melanocytes and nevi.Recent evidence suggests that TRP-1
	is involved in maintaining stability of tyrosinase protein and modulating its catalytic activity.
	TRP-1 is also involved in maintenance of melanosome ultrastructure and affects melanocyte
	proliferation and cell death.

Recombinant human TYRP1 protein

## **Target Details**


Target:	Tyrosinase-Related Protein 1 (TYRP1)
Alternative Name:	TYRP1 (TYRP1 Products)

## **Target Details**

Molecular Weight:	75kDa
Gene ID:	7306
UniProt:	P17643


Application Details	
Application Notes:	Positive Control: SK-MEL-23, SK-MEL-19, SK-MEL-30, SK-MEL-37 cells. Melanoma.
	Known Application: Immunohistochemistry (Formalin-fixed) (5-10 µg/mL for 30 minutes at
	RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM
	EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a
	specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 μg/mL

Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months



## **Immunohistochemistry**

**Image 1.** Formalin-fixed, paraffin-embedded human Melanoma stained with TYRP1 Rabbit Polyclonal Antibody using AEC Chromogen (red).



## **Immunohistochemistry**

**Image 2.** Formalin-fixed, paraffin-embedded human Melanoma stained with TYRP1 Rabbit Polyclonal Antibody using AEC Chromogen (red).