

Datasheet for ABIN6940844  
**anti-Ubiquitin B antibody (AA 1-119)**



[Go to Product page](#)

6 Images

Overview

Quantity:	100 µg
Target:	Ubiquitin B (UBB)
Binding Specificity:	AA 1-119
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Ubiquitin B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Flow Cytometry (FACS), Immunofluorescence (IF), Staining Methods (StM)

Product Details

Immunogen:	Recombinant fragment of human Ubiquitin protein (around aa 1-119) (exact sequence is proprietary)
Clone:	UBB-1748
Isotype:	IgG2c kappa
Purification:	Purified by Protein A/G

Target Details

Target:	Ubiquitin B (UBB)
Alternative Name:	UBB ( <a href="#">UBB Products</a> )

## Target Details

---

**Background:** Ubiquitin is a highly conserved and plays an essential role in the ubiquitin-proteasome pathway. In ubiquitination process, it is first activated by forming a thiol-ester complex with the activation component E1, which is then transferred to ubiquitin-carrier protein E2, followed by to ubiquitin ligase E3 for final delivery to epsilon-NH<sub>2</sub> of the target protein lysine residue. IκB, p53, cdc25A, Bcl-2 etc. are shown as targets of ubiquitin-proteasome process as part of regulation of cell cycle progression, differentiation, cell stress response, and apoptosis. Moreover, ubiquitin have been reported to bind covalently with pathological inclusions which are resistant to degradation e.g. neurofibrillary tangles/paired helical filaments in Alzheimer's disease, Lewy bodies seen in Parkinson's disease, and Pick bodies found in Pick's disease etc.

---

**Molecular Weight:** 9kDa

---

**Gene ID:** 7314

---

**UniProt:** [P62979](#), [P62987](#), [P62988](#)

---

**Pathways:** [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Activation of Innate immune Response](#), [Mitotic G1-G1/S Phases](#), [DNA Replication](#), [Toll-Like Receptors Cascades](#), [Synthesis of DNA](#), [Autophagy](#), [EGFR Downregulation](#), [Ubiquitin Proteasome Pathway](#)

## Application Details

---

**Application Notes:** Positive Control: MCF-7, HeLa or Jurkat cells. Alzheimer's Brain.  
Known Application: Flow Cytometry (1-2 µg/million cells), Western Blot (1-2 µg/mL), Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

---

**Restrictions:** For Research Use only

## Handling

---

**Concentration:** 200 µg/mL

---

**Buffer:** 10 mM PBS with 0.05 % BSA & 0.05 % azide.

---

**Preservative:** Sodium azide

---

**Precaution of Use:** This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---



## Handling

Storage: 4 °C, -80 °C

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months

## Images




### Protein Array

**Image 1.** Analysis of Protein Array containing more than 19,000 full-length human proteins using | Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD s) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD s) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

### Flow Cytometry

**Image 2.** Flow Cytometric Analysis of PFA-fixed MCF-7 cells using Ubiquitin-Monospecific Mouse Monoclonal Antibody (UBB/1748) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Brain stained with Ubiquitin-Monospecific Mouse Monoclonal Antibody (UBB/1748).

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN6940844.