



[Go to Product page](#)

Datasheet for ABIN6940900  
**anti-WT1 antibody (AA 1-181)**

4 Images

Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µg   |
| Target:              | WT1  |
| Binding Specificity: | AA 1-181   |
| Reactivity:          | Human, Mouse, Rat                                  |
| Host:                | Mouse  |
| Clonality:           | Monoclonal   |
| Conjugate:           | This WT1 antibody is un-conjugated                 |
| Application:         | Immunohistochemistry (IHC), Staining Methods (StM) |

Product Details

|              |   |
|--------------|---|
| Immunogen:   | Recombinant protein corresponding to residues 1-181 of human WT1.   |
| Clone:       | 6F-H2   |
| Isotype:     | IgG1 kappa  |
| Specificity: | Recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilm's Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic Wilm's tumors. WT1, a sporadic and familial pediatric kidney tumor, is genetically heterogeneous. Wilm's tumor is associated with mutations of WT1, a zinc-finger transcription factor that is essential for the development of the metanephric kidney and the urogenital system. The WT1 gene is normally expressed in fetal kidney and mesothelium, and its expression has been suggested as a marker for Wilm's tumor and mesothelioma. WT1 protein has been identified in proliferative |

## Product Details

---

mesothelial cells, malignant mesothelioma, ovarian carcinoma, gonadoblastoma, nephroblastoma, and desmoplastic small round cell tumor. Lung adenocarcinomas rarely stain positive with this antibody. WT1 protein expression in mesothelial cells has become a reliable marker for the diagnosis of mesotheliomas.

Purification: Purified by Protein A/G

## Target Details

---

Target: WT1

Alternative Name: WT1 ([WT1 Products](#))

Molecular Weight: 47-55kDa

Gene ID: 7490

UniProt: [P19544](#)

Pathways: [Tube Formation](#)

## Application Details

---

Application Notes: Positive Control: K562 cells. Kidney, Testis, Wilm's Tumor or Mesothelioma.  
Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes) Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

---

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C, -80 °C

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images




SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified Wilm's Tumor Mouse Monoclonal Antibody (6F-H2). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Mesothelioma stained with Wilm's Tumor Mouse Monoclonal Antibody (6F-H2).



Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Testis stained with Wilm's Tumor Mouse Monoclonal Antibody (6F-H2).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6940900.