# antibodies -online.com





## Recombinant anti-WT1 antibody



**Images** 



Go to Product page

_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μg
Target:	WT1
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This WT1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

### **Product Details**

Immunogen:	Recombinant human full-length WT1 protein
Clone:	WT1-1434R
Isotype:	IgG
Specificity:	Recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilm's Tumor (WT1) protein.
	The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking
	exon 2-encoded amino acids, frequently found in subsets of sporadic Wilm's tumors.WT1, a
	sporadic and familial pediatric kidney tumor, is genetically heterogeneous. Wilm's tumor is
	associated with mutations of WT1, a zinc-finger transcription factor that is essential for the
	development of the metanephric kidney and the urogenital system. The WT1 gene is normally
	expressed in fetal kidney and mesothelium, and its expression has been suggested as a marke
	for Wilm's tumor and mesothelioma. WT1 protein has been identified in proliferative

### **Product Details**


1 Todaot Betane		
	mesothelial cells, malignant mesothelioma, ovarian carcinoma, gonadoblastoma,	
	nephroblastoma, and desmoplastic small round cell tumor. Lung adenocarcinomas rarely stair	
	positive with this antibody. WT1 protein expression in mesothelial cells has become a reliable	
	marker for the diagnosis of mesotheliomas.	
Purification:	Purified by Protein A/G	
Target Details		
Target:	WT1	
Alternative Name:	WT1 (WT1 Products)	
Molecular Weight:	47-55kDa	
Gene ID:	7490	
UniProt:	P19544	
Pathways:	Tube Formation	
Application Details		
Application Notes:	Positive Control: K562 cells. Wilm's Tumor, mesothelioma or fetal kidney.	
	Known Application: Immunohistochemistry (Formalin-fixed) (1-2 μg/mL for 30 minutes at	
	RT)Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate	
	Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a	
	specific application should be determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-80 °C	
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody	

is stable for 24 months. Non-hazardous. No MSDS required.

**Expiry Date:** 


24 months

#### **Images**




#### **SDS-PAGE**

**Image 1.** SDS-PAGE Analysis of Purified Wilm's Tumor Rabbit Recombinant Monoclonal Antibody (WT1/1434R).



#### **Immunohistochemistry**

Image 2. Formalin-fixed, paraffin-embedded human
Mesothelioma stained with Wilm's Tumor Rabbit
Recombinant Monoclonal Antibody (WT1/1434R).



### **Immunohistochemistry**

**Image 3.** Formalin-fixed, paraffin-embedded Rat Testis stained with Wilm's Tumor Rabbit Recombinant Monoclonal Antibody (WT1/1434R).