antibodies .- online.com





anti-EPX antibody



Overview

Quantity:	100 μg
Target:	EPX
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This EPX antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:

Clone:	EPO104
Isotype:	IgG1 kappa
Specificity:	Peripheral blood granulocytes are classified into neutrophils, basophils and eosinophils
	according to the staining characteristics of their cytoplasmic granules. Granule proteins are
	released by physiologic and pharmacologic stimuli and play important roles in both normal and
	pathological host immune responses. Eosinophil major basic protein and eosinophil peroxidase
	(EPX) are granule proteins specific to the eosinophil. AHE-1 recognizes human EPX, a granule
	protein specific to eosinophils. It does not cross-react with eosinophil major basic protein,
	elastase, cathepsin G, esterase N, thrombin, plasmin, kallikrein, lactoferrin, or transferrin. This
	MAb stains eosinophils only and does not stain other peripheral blood cells, including platelets,
	neutrophils, monocytes, lymphocytes or red blood cells. Human EPX gene product can form a
	tetramer of two light chains and two heavy chains. Other peroxidase family members include

Human eosinophils from a patient with hypereosinophilic syndrome

Product Details

FIOUUCI Details	
	myeloperoxidase (MPO), lactoperoxidase (LPO), and thyroid peroxidase (TPO).
Purification:	Purified by Protein A/G
Target Details	
Target:	EPX
Alternative Name:	EPX (EPX Products)
Molecular Weight:	70kDa (55kDa-Heavy Chain, 15kDa-Light Chain)
Gene ID:	8288
UniProt:	P11678
Application Details	
Application Notes:	Positive Control: MCF-7 cells. Tonsil.
	Known Application: Flow Cytometry (0.5-1 μ g/million cells), Immunofluorescence (0.5-1 μ
	g/mL), Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only
Handling	
Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
	is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months