

Datasheet for ABIN6941014 anti-CD2 antibody

Images



Overview

Quantity:	100 μg
Target:	CD2
Reactivity:	Human, Cat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD2 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Functional Studies (Func)
Product Details	
Immunogen:	Human thymocytes
Clone:	HuLy-m1
Isotype:	IgG2b kappa
Purification:	Purified by Protein A/G
Target Details	
Target:	CD2
Alternative Name:	CD2 (CD2 Products)
Background:	CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also
	designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-
	specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral

blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a

Target Details

Molecular Weight:

Gene ID:

UniProt:

broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction
between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune
system recognition, thereby facilitating communication between helper T lymphocytes and
antigen-presenting cells, as well as between cytolytic effectors and target cells.
50kDa
914

Application Details

Application Natas:	Positive Control: 293T, MOLT-4 or Jurkat cells. Lymph node and tonsil.
Application Notes:	POSITIVE CONTOL 293 L. MOLLE-4 OF JURKAL CERS LYMDIN HODE AND TONSIL


Known Application: Functional Studies (Order Ab without Azide), ,Flow Cytometry (0.5-1 μ g/million cells), ,Immunofluorescence (0.5-1 μ g/mL), ,Optimal dilution for a specific application should be determined.


Restrictions: For Research Use only


P06729

Handling

Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months







SDS-PAGE

Image 1. SDS-PAGE Analysis Purified CD2 Mouse Monoclonal Antibody (HuLy-m1). Confirmation of Integrity and Purity of Antibody.

Flow Cytometry

Image 2. Flow Cytometric Analysis of MOLT-4 cells using CD2 Mouse Monoclonal Antibody (HuLy-m1) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).