antibodies.com

Datasheet for ABIN6941057 anti-CD5 antibody (AA 269-366)

5 Images



Overview

Quantity:	100 µg
Target:	CD5
Binding Specificity:	AA 269-366
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD5 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunofluorescence (IF), Staining Methods (StM)

Product Details

Immunogen:	Recombinant fragment of human CD5 protein (around aa 269-366) (exact sequence is proprietary)
Clone:	CD5-2418
Isotype:	IgG2c kappa
Specificity:	Recognizes a 67 kDa transmembrane protein, which is identified as CD5. The CD5 antigen is
	found on 95 % of thymocytes and 72 % of peripheral blood lymphocytes. In lymph nodes, the
	main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell marker that also reacts with
	a range of neoplastic B-cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma
	(CLL/SLL), mantle cell lymphoma, and a subset (~10 %) of diffuse large B-cell lymphoma. CD5
	aberrant expression is useful in making a diagnosis of mature T-cell neoplasms. Anti-CD5

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6941057 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

	detection is diagnostic in CLL/SLL within a panel of other B-cell markers, especially one that
	includes anti-CD23. Anti-CD5 is also very useful in differentiating among mature small lymphoid
	cell malignancies. In addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from
	thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.
Purification:	Purified by Protein A/G

Target Details

Target:	CD5
Alternative Name:	CD5 (CD5 Products)
Molecular Weight:	67kDa
Gene ID:	921
UniProt:	P06127

Application Details

Application Notes:	Positive Control: 293T, Ramos or MOLT-4 cells. Tonsil or Lymph Node.	
	Known Application: Flow Cytometry (1-2 μ g/million cells),Immunofluorescence (1-2 μ g/mg),	
	Immunohistochemistry (Formalin-fixed) (1-2 μ g/mL for 30 minutes at RT)(Staining of formalin-	
	fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min	
	followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be	
	determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-80 °C	
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody	
	is stable for 24 months. Non-hazardous. No MSDS required.	


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6941057 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

 . I.	1.
ond	lina
land	

Expiry Date:

24 months


Images



SDS-PAGE

Image 1. SDS-PAGE Analysis


Purified CD5-Monospecific Mouse Monoclonal Antibody (CD5/2418). Confirmation of Purity and Integrity of Antibody.



Protein Array

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using CD5-Monospecific Mouse Monoclonal Antibody (CD5/2418). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Images



Flow Cytometry

Image 3. Flow Cytometric Analysis of PFA-fixed Ramos cells. CD5-Monospecific Mouse Monoclonal Antibody (CD5/2418) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).

Please check the product details page for more images. Overall 5 images are available for ABIN6941057.