antibodies .- online.com







anti-CD19 antibody





Overview

Quantity:	100 μg
Target:	CD19
Reactivity:	Human, Monkey
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD19 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant full-length human CD19 protein
Clone:	CVID3-155
Isotype:	IgG1 kappa
Purification:	Purified by Protein A/G

Target Details

Target:	CD19
Alternative Name:	CD19 (CD19 Products)
Background:	CD19 is a transmembrane glycoprotein that contains two extracellular immunoglobulin-like domains. CD19 is present in both benign and malignant B-cells and is considered to be the
	most reliable surface marker of this lineage over a wide range of maturational stages. In normal
	lymphoid tissue, CD19 is observed in germinal centers, in mantle zone cells, and in scattered

	cells of the inter-follicular areas. Anti-CD19 exhibits an overall immunoreactivity pattern similar
	to those of the antibodies against CD20 and CD22. However, in contrast to CD20, expression of
	CD19 is continuous throughout B-cell development and through terminal differentiation of B-
	cells into plasma cells. Anti-CD19 positivity is seen in the vast majority of B-cell neoplasms
	commonly at a lower intensity than normal B-cell counterparts. Plasma cell neoplasms are
	nearly always negative, as are T-cell neoplasms.
Molecular Weight:	95kDa
Cana ID:	030


Molecular Weight:	95kDa
Gene ID:	930
UniProt:	P15391
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling
	Pathway


Application Details


Application Notes:	Positive Control: Raji cells. Tonsil, Lymph Node or Spleen.
	Known Application: Flow Cytometry (1-2 μg/million cells), ,lmmunofluorescence (1-2 μg/mL),
	,Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

Handling

Concentration:	200 μg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months







Flow Cytometry

Image 1. Flow cytometric analysis of lymphocyte-gated PBMCs. CD19 Mouse Monoclonal Antibody (CVID3/155) labeled with CF568 (orange), unstained cells (gray).

Flow Cytometry

Image 2. Flow Cytometric Analysis of Raji cells. CD19 Mouse Monoclonal Antibody (CVID3/155) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).

Immunofluorescence

Image 3. Immunofluorescence staining of Raji cells. CD19 Mouse Monoclonal Antibody (CVID3/155) followed by goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Reddot (Red).

Please check the product details page for more images. Overall 4 images are available for ABIN6941098.